


Pan Dorset Wound Care Formulary – treat according to highest percentage of tissue type



	Dry	Moist	Wet	Leaking	Saturated
Epithelialising Pink, fragile tissue, promote wound maturation	Softpore Or Atrauman + secondary	Atrauman + secondary Or Comfeel Plus Transparent Or Biatain Silicone (incl Lite)			
Granulating Promote granulation, provide a healthy wound bed		Atrauman + secondary Or UrgoTul + secondary Or UrgoTul Absorb +/- Border Or Comfeel Plus Transparent	KerraCel + secondary Or UrgoTul + secondary Or UrgoTul Absorb Border		
		If chronic wound static and not progressing consider UrgoStart Contact + secondary			
Sloughy Remove slough, promote granulation	ActivHeal Hydrogel + secondary Or Medihoney HCS Or KerraLite Cool +/- Border Or Comfeel Plus Transparent Or Suprasorb X+PHMB + secondary	ActivHeal Hydrogel + secondary Or Medihoney HCS or Tulle + secondary Or KerraLite Cool +/- Border Or Suprasorb X+PHMB + secondary	KerraCel + secondary Or Medihoney Apinate + secondary Or UrgoClean + secondary	KerraCel + secondary Or KerraMax Care Or UrgoClean + secondary	
		Consider Larvae Therapy		Consider Topical Negative Pressure	
Necrotic Debride and/or hydrate (unless hard diabetic eschar)	ActivHeal Hydrogel + secondary Or KerraLite Cool +/- Border Or Prontosan Wound Gel X + secondary Or Flaminal Hydro + secondary Or Suprasorb X+PHMB + secondary	ActivHeal Hydrogel + secondary Or KerraLite Cool +/- Border Or Suprasorb X+PHMB + secondary Or Prontosan Wound Gel X + secondary Or Flaminal Hydro + secondary	KerraCel + secondary Or Flaminal Forte + secondary Or Cutimed Sorbact Gel + secondary	Consider referring to Antimicrobial product information	
		Consider Larvae Therapy			


Secondary Dressings and Absorbency



Pan Dorset Wound Care Formulary – treat according to highest percentage of tissue type

	Dry	Moist	Wet	Leaking	Saturated
Fungating and /or malodorous Manage wound and associated presentation	Cover for patient comfort with non-adherent dressings	Prontosan solution			
		Cutimed Sorbact Swabs + secondary Or UrgoTul SSD+ secondary Or Biatain Silicone (including Lite) Or Assess patients pain, if well managed consider Medihoney Gel sheet + secondary	Cutimed Sorbact Swabs + secondary Or UrgoTul SSD + secondary Or KerraMax Care Or UrgoClean Or Medihoney Tulle sheet + secondary	Cutimed Sorbact Swabs + secondary Or UrgoTul SSD + secondary Or KerraMax Care Or Medihoney Tulle sheet + secondary Or UrgoClean + secondary	
		Clinisorb can be added for malodorous wounds			

Secondary Dressings and Absorbency



- Refer to Moisture Pathway for the management of incontinence associated dermatitis and peri-wound skin barrier preparations
- Refer to the Antimicrobial Product Information for further details on criteria for use and prescribing advisories
- Perform regular wound assessments and document your rationale for dressing selection